

¡STOP! READ THE LABEL BEFORE USING THE PRODUCT

**PRECAUTIONS AND WARNINGS FOR USE:**

**THIS PRODUCT WILL STAIN.** Use only as directed. Avoid product on sidewalks, wooden decks, driveways and clothing. Immediately sweep any product off these surfaces onto treatment area to prevent staining. **IF CONCRETE IS WET WHEN CONTACT OCCURS, STAINING MAY BE UNAVOIDABLE.**

*Do not eat, drink or smoke during the application of this product.*

Keep product of reach of children and pets.

Consumption of this product may make animals sick.

Wash hands after handling this product and before eating food.

Do not ingest this product and avoid contact with eyes and skin.

**FIRST AID:**

In extreme cases this product can produce skin irritation. In case of ingestion, wash the mouth with plenty of water. Seek medical assistance and provide a copy of this label.

**ENVIRONMENTAL PROTECTION MEASURES:**

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden, and sweep any product that lands in the driveway, side-walk, or street, back onto your lawn / garden.

Dispose of product container in accordance with State or local regulations.

Do not throw product packaging or empty product container in soils, rivers, lakes, lagoons or other bodies of water.

**GUARANTEE NOTICE:**

The formulator guarantees the correct composition of the product as long as the user acquires it in its original closed packaging.

Mezfer Crown Inc. does not accept any responsibility for damages resulting from improper storage, inadequate application, nor does Mezfer Crown Inc. accept responsibility for damages and prejudices to crop, people and soils.

**WARNING:**

Over-application or excessive use of boron, or use on crops other than those listed as having a boron response, can result in boron toxicity and may cause serious crop injury.



# GROWTH 8-24-4

## Liquid Inorganic Fertilizer

GUARANTEED ANALYSIS (%)	
TOTAL NITROGEN (N)	8.00
AVAILABLE PHOSPHATE (P <sub>2</sub> O <sub>5</sub> )	24.00
SOLUBLE POTASH (K <sub>2</sub> O)	4.00
BORON (B)	0.02
IRON (Fe)	0.05
MANGANESE (Mn)	0.05
ZINC (Zn)	0.10

Derived from: Urea, Tetraoxphosphoric acid, Potassium hydroxide, 10% Borax, Ferrous sulfate, Manganese sulphate, Zinc sulfate

**NET CONTENTS: 0.264 Gal. / 1L**  
**NET WEIGHT: 2.9 LBS. (1.31 Kg)**

EXPIRATION DATE: 2 years after manufacturing date.

BATCH No. AND MANUFACTURING DATE: Printed on bottle.

**FORMULATED BY:**

MEZCLAS Y FERTILIZANTES S.A DE C.V  
Norte 11, Manzana 1, Lote 1. Cd. Industrial,  
Celaya, Guanajuato, Mexico.

**DISTRIBUTED and GUARANTEED BY:**

MEZFER CROWN INC.  
8415 Datapoint Drive, Suite 750  
San Antonio, Texas 78249

## DIRECTIONS FOR USE

**GENERAL INFORMATION**

MEZFER GROWTH 8-24-4 is an acidic formula of nitrogen, phosphorus and potassium recommended to be applied in stages of high energy requirement, such as the beginning of flowering and fruit development. It is a versatile fertilizer that can be applied to the soil as a conditioner and activator of soil nutrients. In foliar use it is a source of assimilable phosphorus and balanced with macro and micronutrients that activate the development of the plant.

### RECOMMENDATIONS

CROP	RATE/A	APPLICATION INSTRUCTIONS
Vegetables: Chili pepper, Tomato, Solanaceae, Cruciferae, Cucurbits, Spinach, Carrot.	27.4 - 54.7 Fl. Oz./A	Apply 67.6 to 135.2 Fl. Oz in 52 Gals. of water immediately after transplanting at the base of the stem. In foliar application, start 10-15 days after transplanting. Repeat every 15 days.
Melon, Watermelon, Zucchini.	27.4 - 54.7 Fl. Oz./A	Apply during vegetative development at 10-day intervals. Especially apply after cutting in the case of zucchini.
Grains: Corn, Sorghum, Wheat, Sugarcane, Oats, Barley and Pasture.	27.4 - 54.7 Fl. Oz./A	Water the plants starting 20-25 days after emergence. Repeat the application with an interval of 15 days.
Strawberries	54.7 Fl. Oz./A	Apply immediately after transplanting directed to the base of the stem. It can be applied by foliar application in planters and in commercial sowing with application intervals of 15 days.
Potato	27.4 - 136.8 Fl. Oz./A	Apply every 15 days, starting 30 days after the plant is born.
Citrus	54.7 Fl. Oz./A	Apply every 15 days, the critical time is the sprouting of buds, flowering, fruit formation and ripening.
Legumes: Beans, Soy Bean, Broad bean, Lentil and Peanut.	27.4 - 54.7 Fl. Oz./A	Apply every 15 days, starting when the plants have formed 4-5 leaves.
Fruit trees	27.4 - 54.7 Fl. Oz./A	Apply every 15 days, the critical time is the sprouting of buds, flowering, fruit formation and ripening.
Cotton	41.0 - 68.4 Fl. Oz./A	Apply every 15 days, the critical time is the formation of leaves, in flowering and the ripening of acorns.
Ornamental: Rose, Carnation, Crysanthemum, Bird of Paradise, overall cut flowers and nursery.	27.4 - 54.7 Fl. Oz./A	Apply immediately after transplanting directed to the base of the stem.

**METHODS TO PREPARE AND APPLY THE PRODUCT**

Shake the container well before opening. Open with caution avoiding product spillage. Measure the amount of product to apply according to the recommended dose in a volumetric container and add it to the amount of water required according to the type of application and crops. In the case of applications to the neck of the plant and foliar the recommended dose should be divided into 52 Gals./A of water. Do not exceed 67.6 Fl. Oz./ 26 Gals./A of water in foliar application.

**IMPORTANT**

This product has an acid reaction, therefore it **MUST NOT BE MIXED** with alkaline reaction products.

**COMPATIBILITY**

Before doing any mixture, you should consult with your technical advisor and do compatibility testing.

