

¡STOP! READ THE LABEL BEFORE USING THE PRODUCT

PRECAUTIONS AND WARNINGS FOR USE:

THIS PRODUCT WILL STAIN. Use only as directed. Avoid product on sidewalks, wooden decks, driveways and clothing. Immediately sweep any product off these surfaces onto treatment area to prevent staining. **IF CONCRETE IS WET WHEN CONTACT OCCURS, STAINING MAY BE UNAVOIDABLE.**

Do not eat, drink or smoke during the application of this product. Keep out of reach of children and pets.

Consumption of this product may make animals sick. Wash hands after handling this product and before eating food. Do not ingest this product and avoid contact with eyes and skin.

FIRST AID:

In extreme cases this product can produce skin irritation. In case of ingestion, wash the mouth with plenty of water. Seek medical assistance and provide a copy of this label.

ENVIRONMENTAL PROTECTION MEASURES:

Respect good agricultural practices.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn / garden, and sweep any product that lands in the driveway, side-walk, or street, back onto your lawn/garden.

Dispose of product container in accordance with State or local regulations. Do not throw product packaging or empty product container in soils, rivers, lakes, lagoons or other bodies of water.

GUARANTEE NOTICE:

The formulator guarantees the correct composition of the product as long as the user acquires it in its original closed packing. Mezfer Crown Inc. does not accept any responsibility for damages resulting from improper storage, inadequate application, nor does Mezfer Crown Inc. accept responsibility for damages and prejudices to crop, people and soils.

WARNING:

Over-application or excessive use of boron, or use on crops other than those listed as having a boron response, can result in boron toxicity and may cause serious crop injury.



STARTER 15-10-5

Liquid Inorganic Fertilizer

GUARANTEED ANALYSIS (%)	
TOTAL NITROGEN (N)	15.00
AVAILABLE PHOSPHATE (P ₂ O ₅)	10.00
SOLUBLE POTASH (K ₂ O)	5.00
CALCIUM (Ca)	0.05
MAGNESIUM (Mg)	0.05
SULFUR (S)	0.50
BORON (B)	0.02
IRON (Fe)	0.10
MANGANESE (Mn)	0.05
ZINC (Zn)	0.05

Derived from: Urea, Ammonium nitrate, Ammonium polyphosphate, Potassium thiosulfate, 10% Borax, Ferrous sulfate, Manganese sulphate, Magnesium sulphate, Calcium nitrate, Zinc sulfate, Copper sulphate

NET CONTENTS: 0.264 Gal. / 1L
NET WEIGHT: 2.6 Lbs. (1.17 Kg)

EXPIRATION DATE: 2 years after manufacturing date.

BATCH No. AND MANUFACTURING DATE: Printed on bottle.

FORMULATED BY:
MEZCLAS Y FERTILIZANTES S.A DE CV
Norte 11, Manzana 1, Lote 1. Cd. Industrial,
Celaya, Guanajuato, Mexico.

DISTRIBUTED AND GUARANTEED BY:
MEZFER CROWN INC.
8415 Datapoint Drive, Suite 750
San Antonio, Texas 78249

DIRECTIONS FOR USE

GENERAL INFORMATION

MEZFER STARTER 15-10-5 is a fertilizer indicated to favor the growth of the roots, flowers and fruits, recommended to be used in the development stages of the crop. It provides strength and restores crops after periods of stress from drought, flood, or pest and disease damage. The elements it contains act as growth and vigor promoters of plants, both in their initial stage and during advanced stages of crops.

RECOMMENDATIONS

CROP	RATE/A	APPLICATION INSTRUCTIONS
Alfalfa	27.4 - 54.7 Fl. Oz./A	For forage apply 15 days after cutting.
Cotton	27.4 - 54.7 Fl. Oz./A	Apply in the desiccation season, at the time of kiting and when it is in the formation of the cotton.
Coffee tree, Avocado tree.	27.4 - 54.7 Fl. Oz./A	Make applications after flowering one month before cutting and immediately after cutting.
Cabbage, Lettuce	27.4 - 54.7 Fl. Oz./A	Make three applications with intervals of 7-10 days, starting 15 days after transplantation.
Strawberries	27.4 - 54.7 Fl. Oz./A	Apply after transplanting, in flowering and when the fruit is green.
Legumes: Beans, Soy Bean, Haba, Lentil and Peanut.	41.0 - 68.4 Fl. Oz./A	Make the first application 30 days after sowing and apply another 3 at 15-day intervals.
Fruit trees: Apple, Peach, Walnut.	13.53 - 20.29 Fl. Oz.	Make applications after cutting, after flowering and when the fruits are developing.
Tropical fruits: Orange, Lemon, Banana	13.53 - 20.29 Fl. Oz. in 10.69 Gals. water / A	Make applications after cutting, after flowering and when the fruits are developing and make a fourth application after harvest time.
Grains: Corn, Sorghum, Wheat, Sugarcane, Oats, Barley and Pasture.	27.4 - 54.7 Fl. Oz./A	When the plant is between 15-20 cm high at the beginning of the heading and when the grain filling begins, make 2-3 applications.
Transplant vegetables	27.4 - 54.7 Fl. Oz./A	During the growing season, make four applications with an interval of 8 to 10 days.
Direct sowing vegetables	27.4 - 54.7 Fl. Oz./A	Start the applications when the 2nd true leaf appears, making 4 applications with intervals of 8 to 10 days.
Ornamental Rose, Carnation, Crysanthemum, Bird of Paradise, overall cut flowers and nursery.	13.53 - 20.29 Fl. Oz. in 10.69 Gals. water / A	Apply in the first stage of development until the flower buds.
Tobacco	27.4 - 54.7 Fl. Oz./A	Applications are started 15 days after transplantation, subsequently they are carried out every 3 weeks.

METHODS TO PREPARE AND APPLY THE PRODUCT

With the appropriate personal protective equipment on, carefully open the container to avoid spilling the product, mix in 10 to 21 Gals./A of water for land applications and in 15 to 18 Gals./A, of water for aerial applications. The recommendations vary according to the age of the crop and more favorable results are achieved by repeating low doses compared to a single high dose application.

COMPATIBILITY

Before doing any mixing, you should consult with your technical advisor and do compatibility testing.

